

Charge and critical density of strange quark matter

G. X. Peng,^{1,2,3} H. C. Chiang,² P. Z. Ning,³ and B. S. Zou²

¹China Center of Advanced Science and Technology (World Laboratory), Beijing 100080, China

²Institute of High Energy Physics, Academia Sinica, Beijing 100039, China

³Department of Physics, Nankai University, Tianjin 300071, China

(Received 16 November 1998)

The electric charge of strange quark matter is of vital importance to experiments. A recent investigation shows that strangelets are most likely highly negatively charged, rather than slightly positively charged as previously believed. Our present study indicates that negative charges can indeed lower the critical density, and thus be favorable to the experimental searches in heavy ion collisions. However, too much negative charges can make it impossible to maintain flavor equilibrium. {S0556-2813(99)02506-6}

PACS numbers: 24.85.+p, 12.39.-x, 12.38.Mh, 25.75.-q

The last 20 years have witnessed an ever-increasing interest in strange quark matter (SQM) [1]. Early in 1970's [2], it had already been known that the appearance of s fraction in two flavor quark droplets provides an additional degree of freedom, and thus could lower the energy of the system. In 1984, Witten conjectured that quark matter with strangeness per baryon of order unity might be bound [3], which has aroused a great focus of investigations into the stability and detectability of SQM. In the framework of MIT bag model, Jaffe *et al.* find that SQM is absolutely stable around the normal nuclear density for a wide range of parameters [4] while other authors find a few destability factors [5,6]. On application of the quark mass-density-dependent model, Chakrabarty *et al.* carried out a lot of investigations with significantly different results [7,8]. However, the investigations by Benvenuto and Lugones give similar results to those in the bag model [9]. A more recent investigation indicates a link of SQM to the study of quark condensates [10].

The most important way of the territorial searches for SQM is in the ultra-relativistic heavy ion collisions [11]. Of vital importance to the experimental searches is the electric charge of SQM. Previously, it is generally believed that SQM is slightly positively charged [4]. Contrary to previous findings, a recent investigation shows that metastable strangelets are most likely highly negatively charged [12]. This significant result may, if really the case, have a serious impact on the high sensitivity searches in heavy ion experiments at the AGS and CERN facilities. Our present study, in a very simple manner, indicates that proper negative charges can indeed lower the critical density under which SQM can not maintain its flavor equilibrium, and thus no longer exist.

Following Ref. [4], We assume the SQM to be a Fermi gas mixture of u , d , s quarks and electrons with chemical equilibrium maintained by the weak interactions: $d, s \leftrightarrow u + e + \bar{\nu}_e$, $s + u \leftrightarrow u + d$.

At zero temperature, the thermodynamic potential density of the species i is

$$\Omega_i = -\frac{g_i}{48\pi^2} \left[\mu_i(2\mu_i^2 - 5m_i^2)\sqrt{\mu_i^2 - m_i^2} + 3m_i^4 \ln \frac{\mu_i + \sqrt{\mu_i^2 - m_i^2}}{m_i} \right], \quad (1)$$

where $i = u, d, s, e$ (ignore the contribution from neutrinos); g_i is the degeneracy factor with values 6 and 2, respectively for quarks and electrons.

The corresponding particle number density is

$$n_i = -\frac{\partial \Omega_i}{\partial \mu_i} = \frac{g_i}{6\pi^2} (\mu_i^2 - m_i^2)^{3/2}. \quad (2)$$

For a given baryon number density n_b and electric charge density Q , the chemical potentials μ_u , μ_d , μ_s , and μ_e are determined by the following equations

$$\mu_d = \mu_s \equiv \mu, \quad (3)$$

$$\mu_u + \mu_e = \mu, \quad (4)$$

$$\frac{1}{3}(n_u + n_d + n_s) = n_b, \quad (5)$$

$$\frac{2}{3}n_u - \frac{1}{3}n_d - \frac{1}{3}n_s - n_e = Q. \quad (6)$$

The last two equations are equivalent to

$$n_u - n_e = n_b + Q, \quad (7)$$

$$n_d + n_s + n_e = 2n_b - Q. \quad (8)$$

We therefore define a function of μ_e

$$F(\mu_e) = \pi^2(n_d + n_s + n_e) - \pi^2(2n_b - Q) \quad (9)$$

$$= (\mu^2 - m_d^2)^{3/2} + (\mu^2 - m_s^2)^{3/2} + \frac{1}{3}(\mu_e^2 - m_e^2)^{3/2} - \pi^2(2n_b - Q), \quad (10)$$

where

$$\mu = \mu_e + \sqrt{m_u^2 + [\pi^2(n_b + Q) + \frac{1}{3}(\mu_e^2 - m_e^2)^{3/2}]^{2/3}}. \quad (11)$$

Because $m_s > m_{u,d}$, the equation $F(\mu_e) = 0$ for μ_e has solution if and only if

$$\mu \geq m_s, \quad (12)$$

$$F(\mu_e) \leq 0. \quad (13)$$

At the critical density n_c of baryon number, the equality signs in the above two inequalities should be taken. So we can easily find the equation

$$\left\{ m_s - \sqrt{m_u^2 + \left[3\pi^2 n_c - (m_s^2 - m_d^2)^{3/2} \right]^{2/3}} \right\}^2 = m_e^2 + \left\{ 3 \left[\pi^2 (2n_c - Q) - (m_s^2 - m_d^2)^{3/2} \right] \right\}^{2/3} \quad (14)$$

which determines how the critical density depends on charge density, namely, $n_c = f(Q)$.

At n_c , the strangeness fraction becomes zero. When the density decreases further, the equation group (3-6) which determines the configuration of the system has no solution. This indicates that n_c is the lowest density for the possible existence of SQM.

It is obviously difficult to get an explicit expression for the function $f(Q)$. However, the inverse function $f^{-1}(n_c)$ can be easily obtained from Eq. (14):

$$Q = 2n_c - \frac{1}{\pi^2} (m_s^2 - m_d^2)^{3/2} - \frac{1}{3\pi^2} \left[(m_s - \mu')^2 - m_e^2 \right]^{3/2}, \quad (15)$$

where

$$\mu' \equiv \sqrt{m_u^2 + \left[3\pi^2 n_c - (m_s^2 - m_d^2)^{3/2} \right]^{2/3}}. \quad (16)$$

In order to make the inverse square root meaningful, we must require

$$(m_s^2 - m_d^2)^{3/2} \leq 3\pi^2 n_c. \quad (17)$$

When taking the equality sign, and then solving the corresponding equation, we obtain the minimum critical density n_{cmin} . Substituting into Eq. (15) gives

$$Q_{min} = -n_{cmin} - \frac{1}{3\pi^2} \left[(m'_s - m'_u)^2 - m_e^2 \right]^{3/2}, \quad (18)$$

where m'_s and m'_u are, respectively, the masses of s quarks and u quarks corresponding to the density n_{cmin} .

If the charge density Q is more negative than Q_{min} , there is no critical density corresponding to it. Therefore, Q_{min} is the most negative charge density of SQM.

To perform a concrete calculation of the critical density corresponding to $Q > Q_{min}$, we must stop here for a little while to discuss the consideration of quark confinement.

Let's schematically write the QCD Hamiltonian density as

$$H_{QCD} = H_k + \bar{q}M_0q + H_I, \quad (19)$$

where H_k is the kinetic term, H_I is the strong interaction terms between quarks, M_0 is the current mass matrix with diagonal elements m_{u0} , m_{d0} , and m_{s0} .

In the mass-density-dependent model, the strong interaction between quarks is mimicked by the proper dependence of quark masses on density. It is equivalent to expressing the Hamiltonian density as

$$H_{equiv} = H_k + \bar{q}Mq \quad (20)$$

which is the same with that of free quarks except that the current mass matrix M_0 is replaced by an equivalent mass matrix M with diagonal elements m_q ($q = u, d, s$). From the equality

$$\langle n_b | H_{equiv} | n_b \rangle = \langle n_b | H_{QCD} | n_b \rangle, \quad (21)$$

where $|n_b\rangle$ is the state vector with baryon number density n_b , we have

$$m_q = m_{q0} + \frac{\langle n_b | H_I | n_b \rangle}{\sum_q \langle n_b | \bar{q}q | n_b \rangle} \quad (22)$$

$$\equiv m_{q0} + m_I. \quad (23)$$

The physical meaning of Eq. (23) is clear: the first term on the right is the original mass, i.e., the quark current mass, while the second term m_I is the interacting part mimicking the strong interaction between quarks. Obviously, m_I is flavor-independent and density-dependent. Because the characteristic of strong interactions is the confinement of quarks, we should require $\lim_{n_b \rightarrow 0} m_I = \infty$. A popularly used parametrization form is [7-9]

$$m_q = m_{q0} + \frac{B}{3n_b}, \quad (24)$$

where B is a fixed constant determined by stability arguments.

Because we are interested in the true ground state of SQM which might pass into a detector, we must require, at zero pressure, the energy per baryon is less than $M_{56Fe}/56 = 930$ MeV for SQM and greater than 930 MeV for two-flavor quark matter (where M_{56Fe} is the mass of ^{56}Fe) [4]. A little later, we will see that the negative charge density is not permitted to shift too far away from zero. We thus use the same method with that in Ref. [9] to estimate the absolute stability range for the parameter B (the authors of Ref. [9] rename it C), which gives $(69, 110)$ MeV·fm $^{-3}$. Taking the modest values $B = 80$ MeV·fm $^{-3}$, $m_{u0} = m_{d0} = 7.5$ MeV, and $m_{s0} = 100$ MeV, we can calculate the critical density by substituting Eq. (24) into Eq. (14) or (15). The result is plotted in Fig. 1 with a solid line. It is obvious that n_c is an increasing function of Q . Negative charges correspond to smaller critical densities. However, when Q decreases further toward negative direction, there will be no n_c corresponding to it because SQM can not maintain its flavor equilibrium at that case. Therefore, we

conclude that proper negative charges can lower the critical density, and thus be favorable to the experimental searches.

To explore if the above conclusion depends on mass formulas used, we now adopt the following expression:

$$m_q = m_{q0} + \frac{D}{n_b^2}, \quad (25)$$

obtained for reasons of symmetry (particle-antiparticle conjugation, Lorentz transformations). Eq. (25) is obviously in consistency with the properties of the mass (invariant under particle-antiparticle conjugation, Lorentz-scalar; here the density is replaced by square of the four-current $j_\mu j^\mu$). The constant D should be in the range of $(113\text{--}130 \text{ MeV})^7$ according to the method in Ref. [9]. Taking $D = (120 \text{ MeV})^7$ and the same values of m_{q0} with the above, we plot the corresponding critical density as a function of the charge density in Fig. 1 with a dashed line.

We also perform a calculation in the bag model where the quark confinement is mimicked by adding to the energy density expression an extra constant which is interpreted as the vacuum pressure, while the quark masses are density-independent:

$$m_{u,d} = 7.5 \text{ MeV}, \quad (26)$$

$$m_s = 100 \text{ MeV}. \quad (27)$$

The corresponding result is given in Fig. 1 with a dotted line.

The shapes of the three lines in Fig. 1 are obviously similar to each other and so lead to the same conclusion as mentioned in the above. The significant difference is that the line corresponding to bag model calculation is much lower, or in other words, the charge density of stable bulk SQM in the bag model is nearly not allowed to be negative. This is due to the complete ignorance of the strong interactions in the bag model calculation. At high densities, they asymptotically become identical as expected.

It should be pointed out that our method of reasoning is different from what given by Schaffner-Bielich *et al.* [12]. There the investigation was concerned with how small metastable strangelets (including explicit finite size corrections) look like and might decay for different scales of lifetime. None of the candidates found was being absolutely stable, but this fact, of course, depends on the bag parameter being employed. There it was not assumed, as done here, that the strangelets are in weak chemical equilibrium, but it was investigated how the metastable candidates might look like if they are assumed to be stable against strong hadronic decay and subsequently against weak hadronic decay. These decay modes drive the more stable candidates to negative charges. Here we have assumed that the pieces of SQM are in perfect weak chemical equilibrium and considered as having the possibility of absolute stability. Within the standard MIT approach,

these pieces of stable bulk SQM would have a slightly positive charge. It is shown here, with the quark mass-density-dependent model, that the overall charge in fact might be slightly negative. Naturally, the present treatment is very simple, so the concrete values should not be taken very seriously, and further studies are needed.

In conclusion, the present investigation is in favor of the viewpoint that *proper* negative charges can lower the critical density and consequently be favorable to the experimental searches for strange quark matter in ultra-relativistic heavy ion collisions.

ACKNOWLEDGMENTS

The authors would like to thank the partial financial support by National Natural Science Foundation of China under Grant Nos. 19675044 and 19875026.

-
- [1] G.X. Peng, P.Z. Ning, and S.J. Luo, CCAST-WL Workshop Series, **57**, 1 (1996).
 - [2] A.R. Bodmer, Phys. Rev. D **4**, 1601 (1971); S.A. Chin and A.K. Kerman, Phys. Rev. Lett. **43**, 1292 (1979).
 - [3] E. Witten, Phys. Rev. D **30**, 272 (1984).
 - [4] E. Farhi and R. L. Jaffe, Phys. Rev. D **30**, 2379 (1984); M.S. Berger and R.L. Jaffe, Phys. Rev. C **35**, 213 (1987); E.P. Gilson and R.L. Jaffe, Phys. Rev. Lett. **71**, 332 (1993).
 - [5] Jes Madsen, Phys. Rev. Lett. **61**, 2909 (1993); Phys. Rev. D **47**, 5156 (1993); **50**, 3328 (1994); Jes Madsen, Dan M. Jensen, and Michael B. Christiansen, Phys. Rev. C **53**, 1883 (1996).
 - [6] B.C. Parija, 1993, Phys. Rev. C **48**, 2483 (1993); **51**, 1473 (1995).
 - [7] S. Chakrabarty, S. Raha, and B. Sinha, Phys. Lett. B **229**, 112 (1989).
 - [8] S. Chakrabaty, Phys. Rev. D **43**, 627 (1991); **48**, 1409 (1993); **54**, 1306 (1996).
 - [9] O.G. Benvenuto and G. Lugones, Phys. Rev. D **51**, 1989 (1995); G. Lugones and O.G. Benvenuto, *ibid.* **52**, 1276 (1995).
 - [10] G.X. Peng, P.Z. Ning, and H.C. Chiang, Phys. Rev. C **56**, 491 (1997).
 - [11] C. Greiner, P. Koch, and H. Stöcker, Phys. Rev. Lett. **58**, 1825 (1987); C. Greiner, D. H. Rischke, H. Stöcker, and P. Koch, Phys. Rev. D **38**, 2797 (1988). C. Greiner, P. Koch, and H. Stöcker, *ibid.* **44**, 3517 (1991).
 - [12] Jürgen Schaffner-Bielich, Carsten Greiner, Alexander Diener, and Horst Stöcker, Phys. Rev. D **55**, 3038 (1997).

FIG. 1. Critical density n_c vs electric charge density Q . The three lines correspond to different mass formulae (for details, see text). We can see that n_c is an increasing function of Q , i.e., negative charges can lower the critical density. The points marked with “○” correspond to the most negative charge density permitted.

